

# New Similarity Measures of Simplified Neutrosophic Sets and Their Applications

Chunfang Liu\*

## Abstract

The simplified neutrosophic set (SNS) is a generalization of fuzzy set that is designed for some practical situations in which each element has truth membership function, indeterminacy membership function and falsity membership function. In this paper, we propose a new method to construct similarity measures of single valued neutrosophic sets (SVNSs) and interval valued neutrosophic sets (IVNSs), respectively. Then we prove that the proposed formulas satisfy the axiomatic definition of the similarity measure. At last, we apply them to pattern recognition under the single valued neutrosophic environment and multi-criteria decision-making problems under the interval valued neutrosophic environment. The results show that our methods are effective and reasonable.

## Keywords

Multi-Criteria Decision-Making, Pattern Recognition, Similarity Measure, Simplified Neutrosophic Sets

## 1. Introduction

Zadeh [1] proposed fuzzy set (FS) to solve uncertain information. After that, it has been successfully used in different areas [2-7]. Atanassov [8] extended FS to the intuitionistic fuzzy set (IFS) which is very effective to deal with vagueness of the information. The IFS has the ability to express fuzziness and uncertainty of practical situations which each element concludes membership degree, non-membership degree and hesitation degree and they are presented by exact numbers. The sum of these three functions is number 1. In general, the degrees may be not exact numbers but interval values. Then the theory of IFS was extended to interval-valued IFS [9]. With the development of the theory of FS and its extension, they have been handled many uncertainties in different real-life problems. But, there are many phenomena cannot be dealt by the FS and its extension. For example, when we ask an expert about a complex statement, he or she may not give the exact answer of the problem and say that the possibility that the statement is true is 0.6, that is false is 0.5 and the degree that he or she is not sure is 0.3. This situation cannot be expressed by FS and IFS, and some new set is needed to express the condition. Neutrosophic set (NS) was first proposed by Smarandache [10,11] from philosophical point of view. A NS is a set that each element has truth membership degree, indeterminacy membership degree and falsity membership degree.

※ This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Manuscript received August 7, 2015; first revision July 15, 2016; second revision December 6, 2016; accepted April 16, 2017.

Corresponding Author: Chunfang Liu (liuchunfang1112@163.com)

\* College of Science, Northeast Forestry University, Harbin, China (liuchunfang1112@163.com)

As we known, it was difficult to apply the NS in real science and engineering fields. Wang et al. [12,13] proposed single valued neutrosophic set (SVNS) and interval valued neutrosophic set (IVNS). SVNS and IVNS are the subclasses of simplified neutrosophic set which was proposed by Ye [14] and he presented the operators and relationships of the sets and gave their different properties. Therefore, some new results of SNS appeared [15,16]. Similarity measure is used to estimate the degree of similarity of two sets. The similarity measures have been applied to various areas, such as personnel assessment, ecology, medical diagnosis, psychology and clustering analysis, and so on. Broumi and Smarandache [17] defined several similarity measures based on the Hausdorff distance measures of neutrosophic sets. Majumdar and Samanta [18] presented several similarity measures of SVNSs based on distance measures. Ye [14,19-22] also presented the similarity measures of IVNSs and SVNSs and applied them to multi-attribute decision-making problems. Based on tangent function, Prarnaik and Mondal [23] proposed the weighted fuzzy similarity measure and used it to the area of medical diagnosis. On the basis of existing similarity measures, we propose the cotangent similarity measures of SNSs and the weighted cotangent similarity measures of SNSs. Then we apply the proposed similarity measures to pattern recognition in the framework of SVNS and multi-criteria decision-making in the framework of IVNS. The results show that our proposed similarity measure methods are effective and reasonable.

The remainder of the paper is constructed as follows: Section 2 gives some preliminaries. Section 3 gives new cotangent similarity measures of SVNSs. Section 4 gives new cotangent similarity measures of IVNSs. Section 5 gives the applications of the proposed similarity measures in pattern recognition and multi-criteria decision-making. Section 6 concludes the paper.

## 2. Preliminaries

### 2.1 Neutrosophic Set (NS)

Suppose  $X$  is a universe of discourse, and its generic element is  $x$ . A neutrosophic set  $A$  in  $X$  is expressed by truth-membership function  $T_A(x)$ , indeterminacy-membership function  $I_A(x)$ , and falsity-membership function  $F_A(x)$ , respectively. The functions  $T_A(x)$ ,  $I_A(x)$ ,  $F_A(x)$  in  $X$  are real standard or nonstandard subsets of  $]^{-}\bar{0}, 1^{+}[$ , i.e.,

$$T_A(x) : X \rightarrow ]^{-}\bar{0}, 1^{+}[$$

$$I_A(x) : X \rightarrow ]^{-}\bar{0}, 1^{+}[$$

$$F_A(x) : X \rightarrow ]^{-}\bar{0}, 1^{+}[$$

where  $]^{-}\bar{0}, 1^{+}[$  is the nonstandard unit interval, which is an extension of the standard interval  $[0,1]$ ,  $^{-}\bar{0} = 0 - \varepsilon$ ,  $1^{+} = 1 + \varepsilon$ ,  $\varepsilon > 0$ . Then, the sum of  $T_A(x)$ ,  $I_A(x)$ ,  $F_A(x)$  has no restriction, that is  $^{-}\bar{0} \leq \sup T_A(x) + \sup I_A(x) + \sup F_A(x) \leq 3^{+}$ .

Since it is difficult to apply NSs to practical problems, Ye [20] reduced NSs of non-standard intervals into standard intervals that still remain the operations of NSs. If the functions  $T_A(x)$ ,  $I_A(x)$ ,  $F_A(x)$  are single subintervals/subsets in the real standard  $[0,1]$ . Then, a simplification of the NS  $A$  is denoted by

$$A = \{ \langle x, T_A(x), I_A(x), F_A(x) \mid x \in X \}$$

It is a subclass of the NS and contains SVN and IVNS.

For one thing, the membership functions  $T_A(x), I_A(x), F_A(x)$  in a SNS  $A$  are exact numbers in the real unit interval  $[0,1]$ , and the sum of the three functions satisfies the inequality  $0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$ , For every  $x$  in  $X$ . In this case, the SNS changes to the SVN. Let  $A$  and  $B$  be two SVN,  $A$  is contained in  $B$  and denoted by  $A \subseteq B$ , if and only if

$$T_A(x) \leq T_B(x), I_A(x) \geq I_B(x), F_A(x) \geq F_B(x)$$

for every  $x$  in  $X$ .

For another, the membership functions  $T_A(x), I_A(x), F_A(x)$  in SNS  $A$  are the subunit intervals of the real unit interval  $[0,1]$ . We have that

$$T_A(x) = [\inf T_A(x), \sup T_A(x)]$$

$$I_A(x) = [\inf I_A(x), \sup I_A(x)]$$

$$F_A(x) = [\inf F_A(x), \sup F_A(x)]$$

for any point  $x$  in  $X$ . In this case, the SNS reduces to the IVNS.

For two IVNSs  $A$  and  $B$ ,  $A$  is contained in  $B$ , that is  $A \subseteq B$ , if and only if

$$\inf T_A(x) \leq \inf T_B(x), \sup T_A(x) \leq \sup T_B(x),$$

$$\inf I_A(x) \geq \inf I_B(x), \sup I_A(x) \geq \sup I_B(x),$$

$$\inf F_A(x) \geq \inf F_B(x), \sup F_A(x) \geq \sup F_B(x)$$

for every  $x$  in  $X$ .

## 2.2 Similarity Measure of SNSs

Similarity measure is supposed to depict the similarity of a set of alternatives. We use fuzzy evaluation theory to determine the evaluating criteria and get the similarity measure of each alternative and obtain the best alternative. Let  $A, B, C$  be three SNSs, a real valued function  $S: SNS(X) \times SNS(X) \rightarrow [0,1]$ , if it satisfies the following axiomatic conditions:

(S1)  $0 \leq S(A, B) \leq 1$ ;

(S2)  $S(A, B) = 1$  if and only if  $A = B$ ;

(S3)  $S(A, B) = S(B, A)$ ;

(S4)  $A \subseteq B \subseteq C$ , then  $S(A, C) \leq S(A, B)$  and  $S(A, C) \leq S(B, C)$ .

Then we call  $S$  a similarity measure of SNS( $X$ ).

### 3. Cotangent Similarity Measure of SVNSSs

The cosine similarity measure [14] is based on the inner product of two vectors divided by the product of their models. The similarity measure from [22] is based on the tangent function. We first give these two formulas, and give examples to show these results are unreasonable.

$$C(A, B) = \frac{1}{n} \sum_{j=1}^n \frac{T_A(x_j)T_B(x_j) + I_A(x_j)I_B(x_j) + F_A(x_j)F_B(x_j)}{\sqrt{T_A(x_j)^2 + I_A(x_j)^2 + F_A(x_j)^2} \sqrt{T_B(x_j)^2 + I_B(x_j)^2 + F_B(x_j)^2}} \tag{1}$$

$$T(A, B) = 1 - \frac{1}{n} \sum_{j=1}^n \tan \left[ \frac{\pi}{4} \max \left\{ |T_A(x_j) - T_B(x_j)|, |I_A(x_j) - I_B(x_j)|, |F_A(x_j) - F_B(x_j)| \right\} \right] \tag{2}$$

**Example 1.** Let  $A_1 = \langle 0.4, 0.2, 0.6 \rangle$ ,  $B_1 = \langle 0.2, 0.1, 0.3 \rangle$  be two SVNSSs, we use the Eq. (1) and (2) to calculate similarity measure of  $A_1$  and  $B_1$ .

By computing, we get  $C(A_1, B_1) = 1$ ,  $T(A_1, B_1) = 0.6897$ , the result is quite different from each other. We find the membership functions of the two sets exist some differences, but the result shows that their similarity measure is 1. If the similarity measure is 1, it shows that the sets are almost the same sets, which contradict with our intuition.

**Example 2.** Let  $A_2 = \langle 0.3, 0.2, 0.4 \rangle$ ,  $B_2 = \langle 0.4, 0.2, 0.3 \rangle$  be two SVNSSs, we use the Eq. (1) and (2) to calculate similarity measure of  $A_2$  and  $B_2$ .

By computing, we get  $C(A_2, B_2) = 0.9945$ ,  $T(A_2, B_2) = 0.9958$ , the result is almost the same with each other. We found the membership functions of the two sets exist some differences, but the result shows that their similarity measure is very similar, it shows that the sets are almost the same sets, which contradict with our intuition.

From the above examples, we see that the existing similarity measure of SVNSSs sometimes will not consistent with our idea. Motivated by the existing similarity measures, we propose a cotangent similarity measure as follows.

Let

$$A = \left\{ \langle x_j; T_A(x_j), I_A(x_j), F_A(x_j) \rangle \mid x_j \in X \right\}$$

$$B = \left\{ \langle x_j; T_B(x_j), I_B(x_j), F_B(x_j) \rangle \mid x_j \in X \right\}$$

be two SVNSSs in  $X = \{x_1, x_2, \dots, x_n\}$ . We propose the cotangent similarity measure as follows:

$$S(A, B) = \frac{1}{n} \sum_{i=1}^n \cot \frac{\pi}{4} \left[ 1 + \left( \left( \left| T_A(x_j) - T_B(x_j) \right| \vee \left| I_A(x_j) - I_B(x_j) \right| \vee \left| F_A(x_j) - F_B(x_j) \right| \right) \right) \right] \tag{3}$$

where the symbol “ $\vee$ ” is the maximum operation. Then, we can easily prove Eq. (3) satisfies definition of similarity measure. (S1), (S2), (S3) are easily get. We just prove (S4).

If  $A \subseteq B \subseteq C$ , then

$$T_A(x_j) \leq T_B(x_j) \leq T_C(x_j)$$

$$I_A(x_j) \geq I_B(x_j) \geq I_C(x_j)$$

$$F_A(x_j) \geq F_B(x_j) \geq F_C(x_j)$$

then we have the following inequalities:

$$\begin{aligned} |T_A(x_j) - T_B(x_j)| &\leq |T_A(x_j) - T_C(x_j)|, |T_B(x_j) - T_C(x_j)| \leq |T_A(x_j) - T_C(x_j)| \\ |I_A(x_j) - I_B(x_j)| &\leq |I_A(x_j) - I_C(x_j)|, |I_B(x_j) - I_C(x_j)| \leq |I_A(x_j) - I_C(x_j)| \\ |F_A(x_j) - F_B(x_j)| &\leq |F_A(x_j) - F_C(x_j)|, |F_B(x_j) - F_C(x_j)| \leq |F_A(x_j) - F_C(x_j)| \end{aligned}$$

Combining these inequalities, since the cotangent function is a decreasing function within the interval  $[0, \frac{\pi}{2}]$ . Hence,  $S(A, C) \leq S(A, B)$  and  $S(A, C) \leq S(B, C)$ . Therefore we complete the proof.

Usually, the differences of elements' importance are considered. Thus, we need to take the weight of each element  $x_i (i = 1, 2, \dots, n)$  into account.

In the following, we develop the weighted similarity measure between SNSs. Let  $\omega_i, (i = 1, 2, \dots, n)$  be the weight of each element  $x_i (i = 1, 2, \dots, n)$ ,  $\omega_i \in [0, 1], \sum_{i=1}^n \omega_i = 1$ . Then, the weighted similarity measure is obtained as follows:

$$WS(A, B) = \sum_{i=1}^n w_j \cot \frac{\pi}{4} \left[ 1 + \left( \left| T_A(x_j) - T_B(x_j) \right| \vee \left| I_A(x_j) - I_B(x_j) \right| \vee \left| F_A(x_j) - F_B(x_j) \right| \right) \right] \tag{4}$$

Especially if  $\omega_j = \frac{1}{n}, j = 1, 2, \dots, n$ . Eq. (4) reduces to (3).

Now we use the proposed Eq. (3) to calculate the similarity measure of Example 1, then we get  $S(A_1, B_1) = 0.6128$ .

Similarly, we use the proposed Eq. (3) to calculate the similarity measure of Example 2, then we get  $S(A_2, B_2) = 0.8540$ .

From the two examples, we found that the proposed formula is coincided with our intuition and effective to calculate the similarity measure of SVNNS.

## 4. Cotangent Similarity Measure of IVNSs

With the same method, we present the formula of similarity measures of IVNSs.

Let

$$A = \left\{ \langle x_j; T_A(x_j), I_A(x_j), F_A(x_j) \rangle \mid x_j \in X \right\}$$

$$B = \left\{ \langle x_j; T_B(x_j), I_B(x_j), F_B(x_j) \rangle \mid x_j \in X \right\}$$

be two IVNSs in  $X = \{x_1, x_2, \dots, x_n\}$ . Then we propose the new similarity measure of IVNSs.

$$IS(A, B) = \frac{1}{n} \sum_{i=1}^n \cot \frac{\pi}{4} \left[ 1 + \left( \begin{array}{l} \left| \inf T_A(x_j) - \inf T_B(x_j) \right| \vee \left| \inf I_A(x_j) - \inf I_B(x_j) \right| \\ \left| \inf F_A(x_j) - \inf F_B(x_j) \right| \vee \left| \sup T_A(x_j) - \sup T_B(x_j) \right| \\ \left| \sup I_A(x_j) - \sup I_B(x_j) \right| \vee \left| \sup F_A(x_j) - \sup F_B(x_j) \right| \end{array} \right) \right] \quad (5)$$

where the symbol “ $\vee$ ” is the maximum operator. Then, we can easily prove Eq. (5) satisfies the definition of the similarity measure. Here we omitted.

Let  $\omega_i, (i=1, 2, \dots, n)$  be the weight of each element  $x_i (i=1, 2, \dots, n)$ ,  $\omega_i \in [0, 1]$ ,  $\sum_{i=1}^n \omega_i = 1$ . Then, we have the following weighted similarity measure.

$$WIS(A, B) = \sum_{j=1}^n \omega_j \cot \frac{\pi}{4} \left[ 1 + \left( \begin{array}{l} \left| \inf T_A(x_j) - \inf T_B(x_j) \right| \vee \left| \inf I_A(x_j) - \inf I_B(x_j) \right| \\ \left| \inf F_A(x_j) - \inf F_B(x_j) \right| \vee \left| \sup T_A(x_j) - \sup T_B(x_j) \right| \\ \left| \sup I_A(x_j) - \sup I_B(x_j) \right| \vee \left| \sup F_A(x_j) - \sup F_B(x_j) \right| \end{array} \right) \right] \quad (6)$$

## 5. Applications

SNS is a very suitable tool to process the incomplete, uncertainty and inconsistent information. Similarity measure is an important mathematical tool to deal with pattern recognition, medical diagnosis, clustering analysis and decision-making. In the following part, we will use the proposed similarity measures to pattern recognition under the SVN environment and multi-attribute decision-making under the IVN environment.

### 5.1 Application to Pattern Recognition under the SVN Environment

Here is a pattern recognition problem. Suppose that there are  $m$  patterns and they are expressed by SVNNSs. Suppose  $A_i = \left\{ \langle x_j; T_A(x_j), I_A(x_j), F_A(x_j) \rangle \right\}, (i=1, 2, \dots, m)$  are  $m$  patterns in the feature space  $X = \{x_1, x_2, \dots, x_n\}$ . Let  $B = \left\{ \langle x_j; T_B(x_j), I_B(x_j), F_B(x_j) \rangle \right\}$  be a sample needed to be recognized. Our aim is to classify the pattern  $B$  to one of the patterns  $A_1, A_2, \dots, A_m$  according to the principle of the maximum similarity measures. The bigger the similarity measure of  $A_i$  and  $B$  is, the more similar  $A_i$  and  $B$  is. Now, we give the steps for the pattern recognitions. First, calculate the similarity measure of  $A_i$  and  $B$ , that is  $S(A_i, B), i=1, 2, \dots, m$ . Second, choose the largest one  $S(A_k, B)$  from  $S(A_i, B)$ ,

$i = 1, 2, \dots, m$ . Third, conclude that the sample  $B$  belongs to the pattern  $A_k$ .

In the following, a pattern recognition problem about the classification of building material [24] is used to illustrate effective of the proposed similarity measures. Suppose that there are five classes of building materials which are represented by SVNSs  $A_i$ ,  $i = 1, 2, \dots, 5$ , the feature space is  $X = \{x_1, x_2, \dots, x_5\}$ , and  $B$  is an unknown material. They are listed as follows:

$$\begin{aligned}
 A_1 &= \{ \langle x_1, 0.4, 0.6, 0.0 \rangle, \langle x_2, 0.3, 0.2, 0.5 \rangle, \langle x_3, 0.1, 0.3, 0.7 \rangle, \\
 &\quad \langle x_4, 0.4, 0.3, 0.3 \rangle, \langle x_5, 0.1, 0.2, 0.7 \rangle \} \\
 A_2 &= \{ \langle x_1, 0.7, 0.3, 0.0 \rangle, \langle x_2, 0.2, 0.2, 0.6 \rangle, \langle x_3, 0.0, 0.1, 0.9 \rangle, \\
 &\quad \langle x_4, 0.7, 0.3, 0.0 \rangle, \langle x_5, 0.1, 0.1, 0.8 \rangle \} \\
 A_3 &= \{ \langle x_1, 0.3, 0.4, 0.3 \rangle, \langle x_2, 0.6, 0.3, 0.1 \rangle, \langle x_3, 0.2, 0.1, 0.7 \rangle, \\
 &\quad \langle x_4, 0.2, 0.2, 0.6 \rangle, \langle x_5, 0.1, 0, 0.9 \rangle \} \\
 A_4 &= \{ \langle x_1, 0.1, 0.2, 0.7 \rangle, \langle x_2, 0.4, 0.4, 0.4 \rangle, \langle x_3, 0.8, 0.2, 0 \rangle, \\
 &\quad \langle x_4, 0.2, 0.2, 0.7 \rangle, \langle x_5, 0.2, 0.1, 0.7 \rangle \} \\
 A_5 &= \{ \langle x_1, 0.1, 0.1, 0.8 \rangle, \langle x_2, 0, 0.2, 0.8 \rangle, \langle x_3, 0.2, 0, 0.8 \rangle, \\
 &\quad \langle x_4, 0.2, 0, 0.8 \rangle, \langle x_5, 0.8, 0.1, 0.1 \rangle \} \\
 B &= \{ \langle x_1, 0.8, 0.2, 0.1 \rangle, \langle x_2, 0.6, 0.3, 0.1 \rangle, \langle x_3, 0.2, 0.1, 0.8 \rangle, \\
 &\quad \langle x_4, 0.6, 0.5, 0.1 \rangle, \langle x_5, 0.1, 0.4, 0.6 \rangle \}
 \end{aligned}$$

We use Eq. (3) to calculate the similarity measure of  $A_i$  and  $B$ ,  $i = 1, 2, \dots, 5$

$$\begin{aligned}
 S(A_1, B) &= 0.6397, \quad S(A_2, B) = 0.7150, \quad S(A_3, B) = 0.6383 \\
 S(A_4, B) &= 0.3211, \quad S(A_5, B) = 0.3628
 \end{aligned}$$

Since  $S(A_2, B)$  is the biggest, we conclude that the unknown material  $B$  belongs to  $A_2$ .

In this section, we use the similarity measure to the application of pattern recognition, by the similarity measure method, we classify the unknown material  $B$  to  $A_2$ . Now we use the Method of reference [14] and [22] to show that our result is more feasible.

**Method of reference [14]:** If we use Eq. (1) to calculate the similarity measure, we get

$$\begin{aligned}
 S(A_1, B) &= 0.8692, \quad S(A_2, B) = 0.8661, \quad S(A_3, B) = 0.8206 \\
 S(A_4, B) &= 0.5621, \quad S(A_5, B) = 0.4244
 \end{aligned}$$

Since  $S(A_1, B)$  is the biggest, we get that the unknown material  $B$  belongs to  $A_2$ .

**Method of reference [22]:** If we use Eq. (2) to calculate the similarity measure, we get

$$\begin{aligned}
 T(A_1, B) &= 0.9863, \quad T(A_2, B) = 0.9961, \quad T(A_3, B) = 0.9860 \\
 T(A_4, B) &= 0.9926, \quad T(A_5, B) = 0.9920
 \end{aligned}$$

Since  $T(A_2, B)$  is the biggest, we get that the unknown material  $B$  belongs to  $A_2$ .

**Comparison and analysis:** The result of method of reference [14] is different from ours, In Section 3, Example 1 showed its drawbacks. The result of the method of reference [22] is the same as ours. But the calculation of  $T(A_i, B)$  is very close to each other which showed that the unknown material  $B$  is almost close to all  $A_i$ . The result of our given formula shows that our method is effective and reasonable.

## 5.2 Application to Multi-Criteria Decision-Making under IVN Environment

In this section, we apply the similarity measures to multi-criteria decision-making problems in IVNSs.

Let  $X = \{a_1, a_2, \dots, a_n\}$  be a set of alternatives and  $C = \{c_1, c_2, \dots, c_m\}$  be a set of criteria. Assume that the weight of the criterion  $c_j$  is  $w_j$ ,  $w_j \geq 0$ ,  $\sum_{j=1}^m w_j = 1$ . The characteristic of the alternative  $a_i$  is represented by IVNS.

$$a_i = \left\{ \langle c_j, T_{a_i}(c_j), I_{a_i}(c_j), F_{a_i}(c_j) \rangle \mid c_j \in C \right\}$$

$$T_{a_i}(c_j) = [\inf T_{a_i}(c_j), \sup T_{a_i}(c_j)]$$

$$I_{a_i}(c_j) = [\inf I_{a_i}(c_j), \sup I_{a_i}(c_j)]$$

$$F_{a_i}(c_j) = [\inf F_{a_i}(c_j), \sup F_{a_i}(c_j)]$$

If there is only one element in the IVNS, for the sake of simplicity, the interval neutrosophic set is denoted by the interval valued neutrosophic value (IVNV), we denote it as:

$$\alpha_{ij} = \left\langle [a_{ij}, b_{ij}], [c_{ij}, d_{ij}], [e_{ij}, f_{ij}] \right\rangle$$

In multi-criteria decision-making problems, we suppose the ideal point is existed and use it to help identifying the best alternative in the decision set. Then we construct the ideal point to evaluate the alternatives. Generally speaking, the evaluation criteria can be classified by benefit criteria and cost criteria. If the criterion belongs to benefit criterion, we set the ideal point as below:

$$a^* = \left\langle \left[ \max_i(a_{ij}), \max_i(b_{ij}) \right], \left[ \min_i(c_{ij}), \min_i(d_{ij}) \right], \left[ \min_i(e_{ij}), \min_i(f_{ij}) \right] \right\rangle$$

If the criterion belongs to cost criterion, we set the ideal point as below:

$$a^* = \left\langle \left[ \min_i(a_{ij}), \min_i(b_{ij}) \right], \left[ \max_i(c_{ij}), \max_i(d_{ij}) \right], \left[ \max_i(e_{ij}), \max_i(f_{ij}) \right] \right\rangle$$

We calculate the similarity measure  $S(a_i, a^*)$  by Eq. (5) and (6), the ranking order of the alternatives can be determined and the best one can be chosen as well.



Here an example adapted from [19] is utilized to illustrate the applicability and validity of the proposed similarity measure. There is a financial company that wants to invest four companies and they are considered as the four potential alternatives: (1) car company  $a_1$ ; (2) food company  $a_2$ ; (3) computer company  $a_3$ ; (4) arms company  $a_4$ . The invest company must make a decision making considering the following three criteria:  $C_1$  (risk analysis);  $C_2$  (growth analysis);  $C_3$  (environmental impact analysis). Among the three criteria, both  $C_1$  and  $C_2$  are benefit criteria, and  $C_3$  is a cost criterion. Assume that the criteria weight vector is  $w = [0.35, 0.25, 0.40]^T$ . The four possible alternatives are to be evaluated under these three criteria and are in the form of IVNSs, as shown in the following neutrosophic decision matrix  $D$ .

$$D = \begin{pmatrix} \langle [0.4, 0.5], [0.2, 0.3], [0.3, 0.4] \rangle \langle [0.4, 0.6], [0.1, 0.3], [0.2, 0.4] \rangle \langle [0.7, 0.9], [0.2, 0.3], [0.4, 0.5] \rangle \\ \langle [0.6, 0.7], [0.1, 0.2], [0.2, 0.3] \rangle \langle [0.6, 0.7], [0.1, 0.2], [0.2, 0.3] \rangle \langle [0.3, 0.6], [0.3, 0.5], [0.8, 0.9] \rangle \\ \langle [0.3, 0.6], [0.2, 0.3], [0.3, 0.4] \rangle \langle [0.5, 0.6], [0.2, 0.3], [0.3, 0.4] \rangle \langle [0.4, 0.5], [0.2, 0.4], [0.7, 0.9] \rangle \\ \langle [0.7, 0.8], [0.0, 0.1], [0.1, 0.2] \rangle \langle [0.6, 0.7], [0.1, 0.2], [0.1, 0.3] \rangle \langle [0.6, 0.7], [0.3, 0.4], [0.8, 0.9] \rangle \end{pmatrix}$$

From the interval valued neutrosophic matrix, we get the ideal alternative as below:

$$a^* = \{ \langle [0.7, 0.8], [0.0, 0.1], [0.1, 0.2] \rangle, \langle [0.6, 0.7], [0.1, 0.2], [0.1, 0.3] \rangle \\ \langle [0.3, 0.5], [0.3, 0.5], [0.8, 0.9] \rangle \}$$

According to Eq. (6), we get  $S(a_1, a^*) = 0.6000$ ,  $S(a_2, a^*) = 0.8906$ ,  $S(a_3, a^*) = 0.7016$ ,  $S(a_4, a^*) = 0.8451$ . Then we rank the alternatives as  $a_2 > a_4 > a_3 > a_1$ . Obviously,  $a_2$  is the best alternative.

**Comparison and analysis:** In this section, we have proposed a method to solve the MADM problem expressed with IVNSs. From the above example and comparison with the method in [18], we see that their result is the same as ours. In [19], they use two formulas to calculate the similarity measure which is given by the Hamming and Euclidean distance measures; while our method is based on the cotangent functions which satisfy the definition of similarity measure. We use cotangent functions to calculate the similarity measure and the result shows that our method is effective and reasonable.

## 6. Conclusion

In this paper, we commented on the similarity measure of SVNss and IVNSs. We gave the new cotangent functions to calculate the similarity measures. They all satisfied the definition of similarity measures. Then the weighted cotangent similarity measures were introduced by considering the importance of each element. At last, we applied the similarity measure to pattern recognition under the single valued neutrosophic environment and multi-criteria decision-making under the interval valued neutrosophic environment. The results show that our methods are effective and reasonable.

## Acknowledgement

This work was supported by the Fundamental Research Funds for the Central Universities (No. 2572017CB29) and Harbin Science Technology Innovation Talent Research Fund (No. 2016RQXJ230).

## References

- [1] L. A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, no. 3, pp. 338-353, 1965.
- [2] L. A. Zadeh, "The concept of a linguistic variable and its application to approximate reasoning (I)," *Information Science*, vol. 8, no. 3, pp. 199-249, 1975.
- [3] L. A. Zadeh, "The concept of a linguistic variable and its application to approximate reasoning (II)," *Information Science*, vol. 8, no. 4, pp. 301-357, 1975.
- [4] L. A. Zadeh, "The concept of a linguistic variable and its application to approximate reasoning (III)," *Information Science*, vol. 9, no. 1, pp. 43-80, 1975.
- [5] L. A. Zadeh, "Toward a generalized theory of uncertainty (GTU): an outline," *Information Sciences*, vol. 172, no. 1-2, pp. 1-40, 2005.
- [6] J. Ye, "A multicriteria decision-making method using aggregation operators for simplified neutrosophic set," *Journal of Intelligent & Fuzzy Systems*, vol. 26, no. 5, pp. 2459-2466, 2014.
- [7] C. Liu and Y. Luo, "Correlated aggregation operators for simplified neutrosophic set and their application in multi-attribute group decision making," *Journal of Intelligent & Fuzzy Systems*, vol. 30, no. 3, pp. 1755-1761, 2016.
- [8] K. Atanassov, "Intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol. 20, no. 1, pp. 87-96, 1986.
- [9] K. T. Atanassov and G. Gargov, "Interval valued intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol. 31, no. 3, pp. 343-349, 1989.
- [10] F. Smarandache, *Neutrosophy: Neutrosophic Probability, Set and Logic*. Rehoboth, DE: American Research Press, 1999.
- [11] F. Smarandache, *A Unifying Field in Logics: Neutrosophic Logic: Neutrosophy, Neutrosophic Set and Neutrosophic Probability*. Rehoboth, DE: American Research Press, 2003.
- [12] H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, "Single valued neutrosophic sets," *Multispace Multistructure*, no. 4, pp. 410-413, 2010.
- [13] H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, *Interval Neutrosophic Sets and Logic: Theory and Applications in Computing*. Phoenix, AZ: Hexis, 2005.
- [14] J. Ye, "Vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making," *International Journal of Fuzzy Systems*, vol. 16, no. 2, pp. 204-211, 2014.
- [15] C. Li and S. Luo, "The weighted distance measure based method to neutrosophic multi-attribute group decision making," *Mathematical Problems in Engineering*, vol. 2016, article ID 3145341, 2016.
- [16] P. Liu and L. Shi, "The generalized hybrid weighted average operator based on interval neutrosophic hesitant set and its application to multiple attribute decision making," *Neural Computing and Applications*, vol. 26, no. 2, pp. 457-471, 2015.
- [17] S. Broumi and F. Smarandache, "Several similarity measures of neutrosophic sets," *Neutrosophic Sets Systems*, vol. 1, pp. 54-62, 2013.
- [18] P. Majumdar and S. K. Samanta, "On similarity and entropy of neutrosophic sets," *Journal of Intelligent & Fuzzy Systems*, vol. 26, no. 3, pp. 1245-1252, 2014.
- [19] J. Ye, "Similarity measures between interval neutrosophic sets and their applications in multicriteria decision-making," *Journal of Intelligent & Fuzzy Systems*, vol. 26, no. 1, pp. 165-172, 2014.

- [20] J. Ye, "Multiple attribute group decision-making method with completely unknown weights based on similarity measures under single valued neutrosophic environment," *Journal of Intelligent & Fuzzy Systems*, vol. 27, no. 6, pp. 2927-2935, 2014.
- [21] J. Ye, "Improved cosine similarity measures of simplified neutrosophic sets for medical diagnoses," *Artificial Intelligence in Medicine*, vol. 63, no. 3, pp. 171-179, 2015.
- [22] J. Ye and J. Fu, "Multi-period medical diagnosis method using a single valued neutrosophic similarity measure based on tangent function," *Computer Methods and Programs in Biomedicine*, vol. 123, pp. 143-149, 2016.
- [23] S. Pramanik and K. Mondal, "Weighted fuzzy similarity measure based on tangent function and its application to medical diagnosis," *International Journal of Innovative Research in Science, Engineering and Technology*, vol. 4, no. 2, pp. 158-164, 2015.
- [24] W. Wang and X. Xin, "Distance measure between intuitionistic fuzzy set," *Pattern Recognition Letters*, vol. 26, no. 13, pp. 2063-2069, 2005.



**Chunfang Liu** <https://orcid.org/0000-0003-0396-4115>

She received M.S. degree in School of Harbin Institute of Technology, China, in 2003. She is studying for her doctorate in Harbin Engineering University. She is currently working at College of Science, Northeast Forestry University. Her research interests includes system engineering, intelligent control and fuzzy systems.